

# Reductive subgroups of a reductive algebraic group over a local field

George McNinch

2021-11-20

## 1 Overview

- This talk concerns recent work of the speaker McNinch (2021) and McNinch (2020) on reductive groups over a local field.
- Ultimately this work originated from attempts to give a different perspective on construction(s) of (DeBacker 2002).
- These notes will be posted at <https://gmcninch-tufts.github.io/math/> (just google for “McNinch Tufts math” if you’d like to find them...)
- I’d like to thank the organizers of this Special Session on Cohomology, Representation Theory, and Lie Theory for the invitation to speak. Bummer we couldn’t be together in Mobile.
  - I’m going to talk about Lie theory. The questions considered are relevant for (some types of) representation theory. And cohomology is at least playing a back-story...

Nevertheless, I realize that my talk is not exactly at the barycenter of the topics one might have expected in this session, so thanks for your patience!

## 2 Reductive groups and certain subgroups

- Let  $F$  be a field of characteristic  $p \geq 0$ , let  $G$  be a reductive group over  $F$ , and let  $\mu_n$  be the group scheme of  $n$ -th roots of unity, for  $n \geq 2$ .
- Proposition: If  $\phi : \mu_n \rightarrow G$  is a homomorphism, then the image of  $\phi$  is contained in a maximal torus of  $G$ .
- when  $p \mid n$ , note that  $\mu_n$  is not a smooth group scheme. When  $n = p$ , the image of  $\phi$  amounts to  $X \in \text{Lie}(G)$  with  $X^{[p]} = X$ .
- There is a natural notion of equivalence for such homomorphisms; we call the equivalence classes “ $\mu$ -homomorphisms” and denote them as  $\phi : \mu \rightarrow G$ .

## 3 $\mu$ -homomorphisms to a split torus

Proposition: If  $T$  is a split torus over  $F$  with co-character group  $Y = X_*(T)$ , there is a bijection  $\bar{x} \mapsto \phi_{\bar{x}}$

$$Y \otimes \mathbb{Q}/\mathbb{Z} = V/Y \xrightarrow{\sim} \{\mu\text{-homomorphisms } \mu \rightarrow T\}$$

where  $x \in V = Y \otimes \mathbb{Q}$ .

## 4 sub-systems and sub-groups

- let  $\phi : \mu \rightarrow G$  be a  $\mu$ -homomorphism with image in a split torus  $T$ , corresponding to the class of  $x \in Y \otimes \mathbb{Q} = V$  in  $V/Y$ .
- the centralizer  $C_G^0(\phi)$  of the image of  $\phi$  is a subsystem subgroup of  $G$
- if  $G$  is split and  $T$  a maximal split torus, and if  $\Phi$  denotes the roots of  $G$  in  $X^*(T)$ , the root system of  $C_G^0(\phi)$  is given by  $\Phi_x = \{\alpha \in \Phi \mid \langle \alpha, x \rangle \in \mathbb{Z}\}$ .
- $\Phi_x$  is the root subsystem determined by the Borel-de Siebenthal procedure from the extended Dynkin diagram of  $G$ .
- we refer to the reductive subgroups of  $G$  that arises as connected centralizers of homomorphisms  $\phi : \mu \rightarrow G$  as subgroups of type  $C(\mu)$ .

## 5 Local fields

- Let  $K$  be a local field, by which I mean the field of fractions of a complete DVR  $\mathcal{A}$
- write  $k = \mathcal{A}/\pi\mathcal{A}$  for the residue field.
- e.g.  $\mathcal{A}$  could be the completion of the ring of integers  $\mathcal{O}_L$  of a number field  $L$  at some non-zero prime ideal  $\mathfrak{p}$ .

Then  $[K : \mathbb{Q}_p] < \infty$  where  $p\mathbb{Z} = \mathbb{Z} \cap \mathfrak{p}$ .

- or  $\mathcal{A}$  could be the completion of the local ring  $\mathcal{O}_X$  where  $X$  is an (smooth, geometrically irreducible) algebraic curve over  $k$ .

Then  $K \simeq \ell((t))$  where  $[\ell : k] < \infty$ .

- we assume throughout that the char. of the residue field  $k$  is  $p > 0$ .

## 6 Reductive groups and splitting fields

- Let  $G$  be a connected and reductive group over the local field  $K$ .
- can always find a finite, separable extension  $K \subset L$  such that  $G_L$  is split.
- Recall that for a finite separable extension  $k \subset \ell$  of the residue field, there is a unique extension – called an unramified extension –  $K \subset L$  for which the “residue field of  $L$ ” is  $\ell$  and  $[L : K] = [\ell : k]$ .
- We suppose that  $(\diamond) : G$  splits over an unramified extension of  $K$  – i.e. that the group  $G_L$  obtained via base-change is split for a suitable unramified extension  $K \subset L$ .

## 7 Unramified groups

- One says that  $(\clubsuit) : G$  is an unramified group over  $K$  if there is a reductive group scheme  $\mathcal{G}$  over  $\mathcal{A}$  for which  $G = \mathcal{G}_K$ .
- Of course, if  $G$  is split over  $K$ , it is a fundamental fact – essentially, the existence theorem for a reductive group scheme over  $\mathcal{A}$  corresponding to a given root datum – that there is a split reductive “Chevalley group scheme”  $\mathcal{G}$  over  $\mathcal{A}$  with  $G = \mathcal{G}_K$ .
- Any unramified group splits over an unramified extension – i.e.  $(\clubsuit) \implies (\diamond)$  – but the converse is not true in general.

## 8 Parahoric group schemes

- The parahoric group schemes attached to  $G$  are certain affine, smooth group schemes  $\mathcal{P}$  over  $\mathcal{A}$  having generic fiber  $\mathcal{P}_K = G$ .
- We just said that  $G$  is unramified over  $K$  if there is a reductive group scheme  $\mathcal{G}$  over  $\mathcal{A}$  with  $G = \mathcal{G}_K$ . Such a group scheme  $\mathcal{G}$  is a parahoric group scheme.
- But in general, parahoric group schemes  $\mathcal{P}$  are not reductive over  $\mathcal{A}$ , even for split  $G$ . In particular, the special fiber  $\mathcal{P}_k$  need not be a reductive group over the residue field  $k$ .

## 9 Levi factors of the special fiber of a parahoric

Suppose that  $G$  splits over an unramified extension of  $K$ , and let  $\mathcal{P}$  a parahoric attached to  $G$ .

...

Studied Levi decompositions of  $\mathcal{P}_k$  in (McNinch 2010), (McNinch 2014), (McNinch 2020).

...

Theorem (McNinch 2020) There is a reductive subgroup scheme  $\mathcal{M} \subset \mathcal{P}$  such that:

- $\mathcal{M}_K$  is a reductive subgroup of  $G$  of type  $C(\mu)$ , and
- $\mathcal{M}_k$  is a Levi factor of the special fiber  $\mathcal{P}_k$ .

...

Remarks:

- note that  $R_u \mathcal{P}_k$  is defined and split over  $k$ , even if  $k$  is imperfect. (Thus  $\mathcal{P}_k$  has a Levi decomposition over  $k$ ).
- parahorics are determined up to  $G(K)$ -conjugacy by  $x \in V = Y \otimes \mathbb{Q}$ , and  $\mathcal{M}_K$  is the centralizer of  $\phi_{\bar{x}}$ . Here  $Y = X_*(S)$  for a max'l split torus  $S$  in  $G$ .

## 10 Main result on nilpotent elements

- Let  $G$  be a reductive group over the local field  $K$ , and suppose that  $G$  splits over an unramified extension.
- Write  $p$  for the char. of the residue field  $k$  of  $K$ , and s'pose  $p > 2h - 2$  where  $h = h(G)$  is the Coxeter number of  $G$  (i.e. the sup of the Coxeter numbers of simple components of  $G_{\bar{K}}$ .)
- Let  $X \in \text{Lie}(G)$  be a nilpotent element.

...

Theorem: (McNinch 2021) There is a  $K$ -subgroup  $M \subset G$  such that:

- $M$  is a reductive subgp of type  $C(\mu)$  containing a maximal  $K$ -torus of  $G$  which is unramified.
- $M$  is an unramified reductive group over  $K$
- $X \in \text{Lie}(M) \subset \text{Lie}(G)$  and  $X$  is geometrically distinguished for  $M$ .

## 11 Primary tool

- let  $G$  be reductive over  $K$ , suppose that  $G$  splits over unramif. ext, and let  $\mathcal{P}$  be a parahoric for  $G$ .
- Choose reductive subgroup scheme  $\mathcal{M} \subset \mathcal{P}$  as in earlier Theorem – thus  $\mathcal{M}_k$  is a Levi factor of  $\mathcal{P}_k$ .
- Suppose that  $p = \text{char}(k) > 2h - 2$  as before.

...

Theorem: (McNinch 2021) Let  $X_0 \in \text{Lie}(\mathcal{P}_k/R_u\mathcal{P}_k) = \text{Lie}(\mathcal{M}_k)$  be nilpotent.

- a. there is a nilpotent section  $\mathcal{X} \in \text{Lie}(\mathcal{M})$  lifting  $X_0$  which is balanced for  $\mathcal{M}$  – i.e.  $C_{\mathcal{M}_k}(\mathcal{X}_k = X_0)$  and  $C_{\mathcal{M}_k}(\mathcal{X}_K)$  are smooth of the same dimension.
- b. Moreover,  $\mathcal{X}$  is balanced for  $\mathcal{P}$  – i.e. the centralizers  $C_{\mathcal{P}_k}(\mathcal{X}_k)$  and  $C_{\mathcal{P}_k}(\mathcal{X}_K)$  are smooth of the same dimension.

...

- I view this as an alternative version of the lifting Theorem of (DeBacker 2002).

Remarks:

- The Main Theorem above is deduced from the Primary Tool in part via the observation that any nilpotent  $X$  may be placed in  $\text{Lie}(\mathcal{M}) \subset \text{Lie}(\mathcal{P})$  for some parahoric  $\mathcal{P}$ .
- in order to control e.g. the dimensions of the centralizers of  $\mathcal{X}_k$  and  $\mathcal{X}_K$ , we actually place  $\mathcal{X}$  in the image of an  $\mathcal{A}$ -homomorphism  $\text{SL}_{2/\mathcal{A}} \rightarrow \mathcal{M}$  and use the representation theory of  $\text{SL}_2$  (which is well-behaved since  $p > 2h - 2$ ).

The techniques used for this construction build on earlier work of McNinch (2005) on optimal  $\text{SL}_2$ -homomorphisms.

## Bibliography

- DeBacker, Stephen. 2002. “Parametrizing Nilpotent Orbits via Bruhat-Tits Theory.” *Annals of Mathematics. Second Series* 156 (1): 295–332. <https://doi.org/10.2307/3597191>.
- McNinch, George. 2005. “Optimal  $\text{SL}(2)$  Homomorphisms.” *Commentarii Mathematici Helvetici. A Journal of the Swiss Mathematical Society* 80 (2): 391–426. <https://doi.org/10.4171/CMH/19>.
- . 2010. “Levi Decompositions of a Linear Algebraic Group.” *Transformation Groups* 15 (4): 937–64. <https://doi.org/10.1007/s00031-010-9111-8>.
- . 2014. “Levi Factors of the Special Fiber of a Parahoric Group Scheme and Tame Ramification.” *Algebras and Representation Theory* 17 (2): 469–79. <https://doi.org/10.1007/s10468-013-9404-4>.
- . 2020. “Reductive Subgroup Schemes of a Parahoric Group Scheme.” *Transformation Groups* 25 (1): 217–49. <https://doi.org/10.1007/s00031-018-9508-3>.
- . 2021. “Nilpotent Elements and Reductive Subgroups over a Local Field.” *Algebras and Representation Theory* 24: 1479–1522. <https://doi.org/10.1007/s10468-020-10000-2>.